

**Supplemental Table C:** Medications taken by participants and included in GEE analyses as time-varying covariates, organized by outcome measures.

**HYPERTENSION MEDICATIONS: Used in analyses of diastolic and systolic blood pressure.**

<b>Medication (generic name)</b>	<b>Drug Class</b>
LISINOPRIL	ace inhibitor
LISINOPRIL/HCTZ	ace inhibitor
DOXAZOSIN	alpha blocker
PRAZOSIN	alpha blocker
LOSARTAN	angiotensin II receptor antagonist
CLONIDINE	antihypertensive
METOPROLOL	beta blocker
PROPRANOLOL	beta blocker
LABETALOL	beta blocker
ATENOLOL	beta blocker
AMLODIPINE	calcium channel blocker
DILTIAZEM	calcium channel blocker
VERAPAMIL	calcium channel blocker
FUROSEMIDE	diuretic
TRIAMTERENE/HYDROCHLOROTHIAZIDE	diuretic
TRIAMTERENE	diuretic
HYDROCHLOROTHIAZIDE	thiazide
CHLORTHALIDONE	thiazide

**DIABETES MEDICATIONS: Used in analyses of fasting glucose, fasting insulin, Framingham Diabetes Risk Score, and HOMA-IR index.**

<b>Medication (generic name)</b>	<b>Drug Class</b>
EXENATIDE	antidiabetic
INSULIN	antidiabetic
INSULIN ISOPHANE	antidiabetic
INSULIN GLARGINE	antidiabetic
INSULIN LISPRO	antidiabetic
METFORMIN	hypoglycemic
GLIPIZIDE	hypoglycemic
GLIMEPIRIDE	hypoglycemic
GLYBURIDE	hypoglycemic

**CHOLESTEROL MEDICATIONS: Used in analyses of HDL and LDL Cholesterol.**

<b>Medication (generic name)</b>	<b>Drug Class</b>
EZETIMIBE	antihyperlipidemic

CHOLESTYRAMINE	bile acid sequestrant
COLESTIPOL	bile acid sequestrant
PRAVASTATIN	statin
LOVASTATIN	statin
SIMVASTATIN	statin
ATORVASTATIN	statin
ROSUVASTATIN	statin

**CHOLESTEROL AND TRIGLYCERIDE LOWERING MEDICATIONS: Used in analyses of HDL and LDL cholesterol and analyses of triglycerides.**

<b>Medication (generic name)</b>	<b>Drug Class</b>
FENOFIBRATE	fibrate
GEMFIBROZIL	fibrate
NIACIN	nicotinic acid