

TABLES FOR ONLINE DATA SUPPLEMENT

Supplement Table 1. Change in odds of offending and service use outcomes after MAT initiation versus other outpatient substance abuse treatment, schizophrenia sub-sample (n=1,554)

| | Jail | | | Any arrest | | | Felony arrest | | | Substance abuse hospitalization | | | Psychiatric hospitalization | | | Emergency department/Crisis | | | Psychiatric Medication Possession Ratio | | |
|---------------------------------------|------|--------------|--------|------------|--------------|-------|---------------|--------------|-------|---------------------------------|--------------|--------|-----------------------------|--------------|--------|-----------------------------|--------------|--------|---|--------------|-------|
| | O.R. | 95% C.I. | p | O.R. | 95% C.I. | p | O.R. | 95% C.I. | p | O.R. | 95% C.I. | p | O.R. | 95% C.I. | p | O.R. | 95% C.I. | p | O.R. | 95% C.I. | p |
| Study group X time-period interaction | 1.38 | (1.04- 1.83) | 0.024 | 0.91 | (0.69- 1.20) | 0.495 | 1.14 | (0.59- 2.18) | 0.698 | 0.88 | (0.58- 1.33) | 0.549 | 0.69 | (0.48- 0.98) | 0.038 | 0.80 | (0.65- 0.98) | 0.030 | 1.46 | (1.10- 1.95) | 0.009 |
| Main effects | | | | | | | | | | | | | | | | | | | | | |
| Study group‡ | 0.68 | (0.53- 0.89) | 0.004 | 0.91 | (0.73- 1.13) | 0.406 | 0.62 | (0.38- 1.02) | 0.059 | 3.02 | (2.19- 4.16) | <.0001 | 1.75 | (1.31- 2.34) | <.0001 | 1.97 | (1.61- 2.41) | <.0001 | 0.63 | (0.49- 0.82) | 0.001 |
| Time period‡ | 0.58 | (0.49- 0.68) | <.0001 | 1.07 | (0.86- 1.33) | 0.572 | 1.90 | (1.21- 2.99) | 0.006 | 0.26 | (0.16- 0.45) | <.0001 | 0.45 | (0.34- 0.59) | <.0001 | 0.71 | (0.58- 0.88) | 0.001 | 0.85 | (0.70- 1.05) | 0.128 |

‡Models control for age, educational need, gender, primary psychiatric diagnosis, race/ethnicity, insurance (SSI, Medicaid), probation, service participation (residential, Assertive Community Treatment, outpatient mental health treatment, outpatient substance abuse treatment), medication possession ratio (prior month medication possession ratio of 80% or higher), time, and community tenure. Medication possession ratio outcome model excludes medication possession ratio covariate, includes only person-months when individuals were enrolled in Medicaid, and is restricted to individuals with any record of medication who were diagnosed with schizophrenia spectrum or bipolar disorder (n=1,035).

‡ MAT group = 1, COMP group = 0; Post-index-treatment period = 1, pre-index-treatment period = 0

Supplement Table 2. Change in odds of offending and service use outcomes after MAT initiation versus other outpatient substance abuse treatment, bipolar disorder sub-sample (n=1,551)

| | Jail | | | Any arrest | | | Felony arrest | | | Substance abuse hospitalization | | | Psychiatric hospitalization 95% | | | Emergency department/Crisis 95% | | | Psychiatric Medication Possession Ratio 95% | | |
|---------------------------------------|------|--------------|--------|------------|--------------|-------|---------------|--------------|-------|---------------------------------|--------------|--------|---------------------------------|--------------|--------|---------------------------------|--------------|--------|---|--------------|-------|
| | O.R. | 95% C.I. | p | O.R. | 95% C.I. | p | O.R. | 95% C.I. | p | O.R. | 95% C.I. | p | O.R. | C.I. | p | O.R. | C.I. | p | O.R. | C.I. | p |
| Study group X time-period interaction | 1.31 | (0.94- 1.81) | 0.107 | 0.98 | (0.75- 1.26) | 0.854 | 0.50 | (0.29- 0.86) | 0.012 | 0.98 | (0.68- 1.41) | 0.917 | 0.78 | (0.53- 1.15) | 0.212 | 0.93 | (0.72- 1.21) | 0.610 | 1.65 | (1.21- 2.24) | 0.002 |
| Main effects | | | | | | | | | | | | | | | | | | | | | |
| Study group‡ | 0.68 | (0.51- 0.90) | 0.007 | 0.96 | (0.79- 1.17) | 0.697 | 1.32 | (0.92- 1.88) | 0.133 | 2.40 | (1.81- 3.17) | <.0001 | 1.75 | (1.30- 2.35) | <.0001 | 1.59 | (1.27- 1.99) | <.0001 | 0.86 | (0.65- 1.15) | 0.313 |
| Time period‡ | 0.48 | (0.41- 0.56) | <.0001 | 0.82 | (0.65- 1.03) | 0.094 | 1.73 | (1.11- 2.70) | 0.016 | 0.14 | (0.09- 0.23) | <.0001 | 0.23 | (0.15- 0.35) | <.0001 | 0.37 | (0.29- 0.48) | <.0001 | 0.90 | (0.70- 1.15) | 0.398 |

‡Models control for age, educational need, gender, primary psychiatric diagnosis, race/ethnicity, insurance (SSI, Medicaid), probation, service participation (residential, Assertive Community Treatment, outpatient mental health treatment, outpatient substance abuse treatment), medication possession ratio (prior month medication possession ratio of 80% or higher), time, and community tenure. Medication possession ratio outcome model excludes medication possession ratio covariate, includes only person-months when individuals

‡ MAT group = 1, COMP group = 0; Post-index-treatment period = 1, pre-index-treatment period = 0

Supplement Table 3. Change in odds of offending and service use outcomes after MAT initiation versus other outpatient substance abuse treatment, by medication possession ratio status and primary psychiatric disorder

| | Jail | | | Any arrest | | | Any felony arrest | | | Substance abuse hospitalization | | | Psychiatric hospitalization | | | Emergency department/Crisis | | |
|--|------|--------------|--------|------------|--------------|-------|-------------------|--------------|-------|---------------------------------|--------------|--------|-----------------------------|--------------|--------|-----------------------------|--------------|--------|
| | O.R. | 95% C.I. | p | O.R. | 95% C.I. | p | O.R. | 95% C.I. | p | O.R. | 95% C.I. | p | O.R. | 95% C.I. | p | O.R. | 95% C.I. | p |
| Sample: Schizophrenia/HIGH Medication Possession Ratio (n=405) | | | | | | | | | | | | | | | | | | |
| Study group X time-period interaction | 0.95 | (0.44- 2.04) | 0.900 | 0.71 | (0.36- 1.40) | 0.323 | 1.04 | (0.24- 4.45) | 0.956 | 0.57 | (0.19- 1.69) | 0.313 | 0.49 | (0.27- 0.91) | 0.024 | 0.91 | (0.61- 1.34) | 0.621 |
| Main effects | | | | | | | | | | | | | | | | | | |
| Study group¥ | 0.61 | (0.38- 0.98) | 0.042 | 1.12 | (0.75- 1.68) | 0.592 | 0.57 | (0.21- 1.59) | 0.283 | 3.06 | (1.67- 5.61) | 0.000 | 2.49 | (1.55- 4.01) | 0.000 | 2.02 | (1.41- 2.89) | 0.000 |
| Time period¥ | 0.66 | (0.45- 0.96) | 0.029 | 0.83 | (0.48- 1.46) | 0.524 | 2.08 | (0.75- 5.74) | 0.159 | 0.13 | (0.03- 0.56) | 0.006 | 0.68 | (0.42- 1.10) | 0.118 | 0.88 | (0.58- 1.32) | 0.524 |
| Sample: Schizophrenia/LOW Medication Possession Ratio (n=564) | | | | | | | | | | | | | | | | | | |
| Study group X time-period interaction | 1.64 | (1.13- 2.37) | 0.009 | 1.05 | (0.70- 1.56) | 0.828 | 1.39 | (0.56- 3.48) | 0.477 | 0.92 | (0.48- 1.75) | 0.799 | 0.96 | (0.58- 1.56) | 0.855 | 0.89 | (0.67- 1.19) | 0.439 |
| Main effects | | | | | | | | | | | | | | | | | | |
| Study group¥ | 0.65 | (0.45- 0.93) | 0.019 | 0.83 | (0.59- 1.18) | 0.301 | 0.57 | (0.26- 1.22) | 0.146 | 3.72 | (2.10- 6.60) | <.0001 | 1.01 | (0.66- 1.53) | 0.979 | 1.94 | (1.44- 2.60) | <.0001 |
| Time period¥ | 0.44 | (0.34- 0.57) | <.0001 | 1.07 | (0.76- 1.51) | 0.685 | 1.21 | (0.59- 2.46) | 0.608 | 0.38 | (0.17- 0.86) | 0.021 | 0.38 | (0.26- 0.56) | <.0001 | 0.72 | (0.53- 0.98) | 0.037 |
| Sample: Bipolar Disorder/HIGH Medication Possession Ratio (n=374) | | | | | | | | | | | | | | | | | | |
| Study group X time-period interaction | 0.87 | (0.47- 1.60) | 0.653 | 0.70 | (0.40- 1.20) | 0.189 | 0.18 | (0.05- 0.64) | 0.008 | 0.66 | (0.34- 1.28) | 0.219 | 0.43 | (0.24- 0.78) | 0.005 | 0.60 | (0.39- 0.91) | 0.016 |
| Main effects | | | | | | | | | | | | | | | | | | |
| Study group¥ | 0.67 | (0.41- 1.07) | 0.091 | 0.98 | (0.67- 1.43) | 0.908 | 2.14 | (1.08- 4.22) | 0.029 | 2.18 | (1.33- 3.56) | 0.002 | 1.71 | (1.05- 2.78) | 0.031 | 1.47 | (1.03- 2.11) | 0.036 |
| Time period¥ | 0.44 | (0.31- 0.64) | <.0001 | 0.76 | (0.47- 1.23) | 0.265 | 1.99 | (0.76- 5.18) | 0.159 | 0.16 | (0.07- 0.39) | <.0001 | 0.46 | (0.22- 0.97) | 0.042 | 0.55 | (0.33- 0.92) | 0.022 |
| Sample: Bipolar Disorder/LOW Medication Possession Ratio (n=532) | | | | | | | | | | | | | | | | | | |
| Study group X time-period interaction | 1.65 | (0.93- 2.92) | 0.084 | 1.12 | (0.75- 1.69) | 0.575 | 0.94 | (0.38- 2.33) | 0.894 | 0.76 | (0.43- 1.34) | 0.347 | 1.25 | (0.68- 2.29) | 0.480 | 1.05 | (0.70- 1.56) | 0.820 |
| Main effects | | | | | | | | | | | | | | | | | | |
| Study group¥ | 0.66 | (0.43- 1.02) | 0.064 | 0.93 | (0.69- 1.26) | 0.637 | 0.93 | (0.51- 1.70) | 0.819 | 3.10 | (2.01- 4.79) | <.0001 | 1.66 | (1.03- 2.68) | 0.036 | 1.72 | (1.18- 2.50) | 0.005 |
| Time period¥ | 0.38 | (0.29- 0.51) | <.0001 | 0.65 | (0.45- 0.95) | 0.024 | 1.00 | (0.48- 2.08) | 0.994 | 0.27 | (0.12- 0.61) | 0.002 | 0.22 | (0.12- 0.40) | <.0001 | 0.36 | (0.25- 0.53) | <.0001 |

‡Models control for age, educational need, gender, race/ethnicity, insurance (SSI, Medicaid), probation, service utilization (residential treatment, Assertive Community Treatment, outpatient mental health treatment, outpatient substance abuse treatment), medication possession ratio, time, and community tenure.

¥ MAT group = 1, COMP group = 0; Post-index-treatment period = 1, pre-index-treatment period = 0

