



FIGURE S1. Pattern of Across-Session Change for Prolonged Exposure and Sertraline


A. Acute Outcome PTSD Severity (PSS-SR): PE versus Sertraline



B. Acute Outcome Depression Severity (BDI): PE versus Sertraline



C. Acute Outcome Anxiety Severity (STAI): PE versus Sertraline



D. Acute Outcome Sheehan Disability Scale (SDS): PE versus Sertraline




TABLE S1. Intent-to-Treat Mean, SD, and Effects Sizes for Treatment Choice versus No Choice

| Choice | Pre-Treatment | | | | Post-Treatment | | | |
|------------------------------------------------|---------------|---------------|-------|--------------|----------------|---------------|------|--------------|
| | <i>M</i> | <i>95% CI</i> | % | (<i>n</i>) | <i>M</i> | <i>95% CI</i> | % | (<i>n</i>) |
| PTSD Symptom Scale-Interview | 29.5 | 29.2-29.8 | | | 10.9 | 9.9-11.9 | | |
| PTSD Symptom Scale-Self Report | 33.6 | 32.0-35.2 | | | 13.4 | 11.3-15.5 | | |
| PTSD Diagnosis | | | 100.0 | (97) | | | 28.9 | (28) |
| Responder Status | | | 0.0 | (0) | | | 73.2 | (71) |
| Hamilton Rating Scale Depression ₂₄ | 24.3 | 23.9-24.7 | | | 12.2 | 11.7-12.7 | | |
| Beck Depression Inventory | 24.0 | 22.1-25.9 | | | 10.0 | 8.2-11.9 | | |
| Spielberger State Anxiety Inventory-State | 56.4 | 54.5-58.3 | | | 40.6 | 38.4-42.8 | | |
| Sheehan Disability Scale | 18.9 | 17.6-20.2 | | | 11.8 | 10.1-13.5 | | |
| No Choice | | | | | | | | |
| | <i>M</i> | <i>95% CI</i> | % | (<i>n</i>) | <i>M</i> | <i>95% CI</i> | % | (<i>n</i>) |
| PTSD Symptom Scale-Interview | 29.7 | 29.4-30.0 | | | 12.5 | 11.2-13.8 | | |
| PTSD Symptom Scale-Self Report | 34.8 | 33.4-36.2 | | | 15.4 | 12.8-18.0 | | |
| PTSD Diagnosis | | | 100.0 | (103) | | | 44.7 | (46) |
| Responder Status | | | 0.00 | (0) | | | 57.3 | (59) |
| Hamilton Rating Scale Depression ₂₄ | 23.1 | 22.7-23.5 | | | 12.6 | 12.1-13.1 | | |
| Beck Depression Inventory | 25.4 | 23.5-27.3 | | | 11.1 | 8.7-13.5 | | |
| Spielberger State Anxiety Inventory-State | 56.5 | 54.4-58.6 | | | 41.9 | 39.1-44.7 | | |
| Sheehan Disability Scale | 18.6 | 17.4-19.8 | | | 10.8 | 9.2-12.4 | | |

Note. Responder Status = PSSI ≤ 24 and CGI < 4; Loss of Diagnosis = Does not meet DSM-IV symptom criteria. Means and 95% CIs were generated from the mixed effect models. Data for outcomes measures are estimated using the intent-to-treat analysis (*N* = 200). For PTSD Diagnosis, $\chi^2(1, N = 200) = 5.35, p = .02$, and for Responder Status, $\chi^2(1, N = 200) = 5.56, p = .02$. All other analyses between Choice and No Choice were not significant.