

**ONLINE ONLY APPENDIX TABLE 1. TREATMENT VISITS FOR BIPOLAR DISORDER STRATIFIED BY DRUG CLASS, 1998-2009.**

	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009
Mood Stabilizers, %	82	79	79	75	72	75	70	72	71	69	68	67
Second generation												
Antipsychotics, %	18	19	25	32	35	37	39	41	45	48	47	49
Antidepressants, %	47	47	48	49	53	52	46	44	42	41	40	41
Anxiolytics, %	21	22	17	14	18	17	17	18	19	18	18	20
Other, %	7	5	4	4	4	4	4	4	4	4	4	5
Total Bipolar Visits (millions)	3.7	3.6	3.6	3.9	5.6	5.9	6.3	7.9	9.1	8.4	8.2	7.8

Column percents exceed 100% because rows are not mutually exclusive (e.g., single visit for bipolar disorder may yield more than one treatment visit if more than one therapy used as treatment); "Other" includes stimulants and typical antipsychotics;  
Source: IMS Health National Disease and Therapeutic Index™, 1998-2009

**ONLINE ONLY APPENDIX FIGURE 1. TREATMENT VISITS FOR BIPOLAR DISORDER DURING WHICH AN SECOND GENERATION ANTIPSYCHOTIC WAS PRESCRIBED, 1998-2009.**

Source: IMS Health National Disease and Therapeutic Index™, 1998-2009



