

Appendix: Mental health and medical comorbidities, active at time of chart review ^a

	Active mental health diagnosis				
	PTSD (n=74)	Mood (n=32)	Anxiety (n=8)	SUD (n=35)	Other MH (n=9)
<i>Active medical comorbidity</i>					
Pain	50 (68%)	24 (75%)	7 (87%)	28 (80%)	8 (89%)
Lung	6 (8%)	2 (6%)	0 (0%)	4 (11%)	0 (0%)
Cardio	46 (62%)	25 (78%)	4 (50%)	21 (60%)	6 (67%)
Diabetes	29 (39%)	18 (56%)	1 (12%)	15 (43%)	3 (33%)
GI	32 (43%)	18 (56%)	4 (50%)	19 (54%)	5 (56%)
Nicotine	15 (20%)	11 (34%)	1 (12%)	12 (34%)	1 (11%)
Other medical	56 (76%)	28 (87%)	5 (62%)	26 (74%)	7 (78%)
No. medical comorbidities: mean (median), range	3.16 (3.0), 0 - 7	3.94 (4.0), 0 - 6	2.75 (3.0), 0 - 4	3.57 (4.0), 0 - 7	3.33 (4.0), 0 - 7
<i>Active mental health comorbidity</i>					
PTSD	—	29 (91%)	6 (75%)	32 (91%)	7 (78%)
Mood	29 (39%)	—	3 (37%)	17 (49%)	4 (44%)
Anxiety	6 (8%)	3 (9%)	—	5 (14%)	2 (22%)
SUD	32 (43%)	17 (53%)	5 (62%)	—	7 (78%)
Other MH	7 (9%)	4 (12%)	2 (25%)	7 (20%)	—

Total no. mental health	2.0	2.66	3.0	2.74	3.22
diagnoses: Mean	(2.0),	(3.0),	(3.0),	(3.0),	(4.0),
(median), range	1 - 4	1 - 4	2 - 3	2 - 4	2 - 4

^a column percents are reported