

Supplementary Table 1. *p*-values for the chi-squared associations between diagnostic variables and number of appointments with receipt of assessment and complete care variables

	Smoking Assessment	Nutrition Assessment	Alcohol Assessment	Physical Activity Assessment	Smoking Complete Care	Nutrition Complete Care	Alcohol Complete Care	Physical Activity Complete Care
Depression	.96	.44	.12*	.82	.29	.67	.85	.76
Bi-polar disorder	.80	.40	.56	.61	.69	.72	.11*	.53
Schizophrenia/other psychotic illness	.51	.63	.13*	.52	.56	.35	.002*	.64
Anxiety disorder	.40	.38	.79	.37	.16*	.98	.26	.98
Other mental illness	.88	.42	.69	.12*	.30	.46	.001*	.43
Heart diseases/stroke	.24*	.69	.17*	.17*	.30	.86	.70	.16*
Diabetes	.66	.05*	.73	.99	.34	.02*	1.00	.005*
Cancers	.66	.59	.51	.45	.35	.02*	.29*	.86
Respiratory	.66	.69	.43	.06*	.77	.03*	.67	.97
Number of appointments in last 12 months	.10*	.06	.71	<.001*	.11*	.36	.003*	.12*

* Variable entered into backwards stepwise regression model