**Efficacy Study #1, Adults
with Severe Mental Illness
Smith et al., 2014**
n=25 VR-JIT trainees
n=12 waitlist controls

**Efficacy Study #2,
Veterans with PTSD
Smith et al., 2015**
n=23 VR-JIT trainees
n=10 waitlist controls

Transition to VR-JIT group
n=5 waitlist control
participants used VR-JIT
after the efficacy study

Transition to VR-JIT group
n=1 waitlist control
participant used VR-JIT
after the efficacy study

**6 Month Follow-Up Study
Eligible Participants**
n=54 VR-JIT
n=16 waitlist controls

Attrition
n=1 deceased
n=1 hospitalized
n=1 retired
n=16 lost contact (including 1 waitlist control
that used VR-JIT)

**6 Month Follow-Up
Completed Participants**
n=39 VR-JIT
n=12 waitlist controls

Supplemental Figure 1. Flow Chart of participants from two efficacy studies evaluated

as a single follow-up cohort

Supplemental Table 1. VR-JIT group status as a predictor of a job offer				
	Step 1		Step 2	
	OR	95% CI	OR	95% CI
Step 1				
Neurocognition	1.06+	1.00-1.13	1.07+	.99-1.15
Months since prior employment	.97**	.95-.99	.96*	.92-.99
PTSD Diagnosis	3.57+	.81-15.85	6.09*	1.08-34.40
Step 2				
VR-JIT group (yes or no)	--	--	9.18*	1.34-62.93

+p<.10, *p<.05, **p<.01;
Step 1 Nagelkerke R^2=.40;
Step 2 Nagelkerke R^2=.54.