

Supplemental Table 1. Current medications and treatment settings of psychiatric participants

	Schizophrenia (N=837)	Bipolar Disorder (N=530)
Psychiatric medications received at time of study visit		
Atypical antipsychotic	696 (83%)	363 (68%)
Clozapine	166 (20%)	5 (1%)
Olanzapine	198 (24%)	70 (13%)
Risperidone	223 (27%)	103 (19%)
Mood stabilizers		
Lithium	104 (12%)	179 (34%)
Valproate	173 (21%)	170 (32%)
Anticonvulsant	346 (41%)	437 (82%)
Antidepressant	347 (41%)	224 (42%)
Treatment setting at time of study visit		
Inpatient	167 (20%)	315 (59%)
Day hospital	55 (7%)	50 (9%)
Outpatient	615 (73%)	165 (31%)

Supplemental Table 2. Number of participants by diagnostic group by year of enrollment

	Schizophrenia	Bipolar Disorder	Non-psychiatric controls
<u>Year</u>			
1999	90	0	0
2000	94	0	0
2001	49	50	0
2002	38	66	71
2003	60	4	28
2004	27	18	0
2005	56	9	66
2006	31	13	35
2007	42	20	45
2008	39	50	65
2009	57	42	61
2010	50	43	36
2011	61	47	49
2012	27	45	36
2013	11	34	37
2014	21	30	12
2015	44	31	13
2016	40	28	17
Total	837	530	571