

Unadjusted racial and ethnic differences in clinical characteristics by treatment condition during 24-month treatment period (Beta's, odd ratios, confidence intervals, p values)

Clinical Characteristics	Community Care								
	Race/Ethnicity; Non-Hispanic white (REF)								
	Non-Hispanic Black			Hispanic			Others		
	β	95% CI	<i>p</i> value	β	95% CI	<i>p</i> value	β	95% CI	<i>p</i> value
PANSS ^a									
Total score	6.42	2.11 to 10.72	.003		NS			NS	
Negative symptoms		NS			NS			NS	
Positive symptoms	3.21	1.36 to 5.07	.001		NS			NS	
Disorganized thought	1.78	.67 to 2.89	.002		NS			NS	
Uncontrolled hostility	.80	.15 to 1.46	.017		NS		-.81	-1.57 to -.04	.039
Anxiety/depression		NS			NS			NS	
Service Utilization	OR	95% CI	<i>p</i> value	OR	95% CI	<i>p</i> value	OR	95% CI	<i>p</i> value
Medication management		NS			NS			NS	
Individual therapy	.45	.28 to .72	.001	.40	.18 to .89	.024		NS	
Family psychoeducation		NS		.18	.05 to .63	.007		NS	
	NAVIGATE								
	Non-Hispanic Black			Hispanic			Others		
	β	95% CI	<i>p</i> value	β	95% CI	<i>p</i> value	β	95% CI	<i>p</i> value
PANSS ^a									
Total score		NS			NS		9.52	.28 to 18.76	.043
Negative symptoms		NS			NS			NS	
Positive symptoms		NS			NS			NS	
Disorganized thought		NS		2.05	.75 to 3.34	.002		NS	
Uncontrolled hostility		NS			NS			NS	
Anxiety/depression	1.28	.01 to 2.55	.48		NS		2.98	.89 to 5.06	.005
Service Utilization	OR	95% CI	<i>p</i> value	OR	95% CI	<i>p</i> value	OR	95% CI	<i>p</i> value
Medication management		NS		2.82	1.68 to 4.74	.001		NS	
Individual therapy		NS			NS			NS	
Family psychoeducation	.49	.30 to .81	.005		NS			NS	

^a Positive and Negative Syndrome Scale. Possible scores range from 30 to 120 with higher scores indicating greater psychosis symptom severity.

^b Baseline Positive and Negative Syndrome Scale subscale score



Unadjusted racial and ethnic differences in clinical characteristics by treatment condition during 24-month treatment period (means and SD)

Clinical Characteristics	Community Care							
	Race/Ethnicity							
	Non-Hispanic white (REF)		Non-Hispanic Black		Hispanic		Others	
	M	SD	M	SD	M	SD	M	SD
PANSS ^a								
Total score	65.13	1.50	71.55	1.61**	65.05	3.52	69.96	7.19
Negative symptoms	17.37	.52	17.93	.62	16.96	1.67	17.09	2.63
Positive symptoms	19.73	.69	22.94	.65***	18.89	1.26	23.76	2.81
Disorganized thought	13.08	.34	14.86	.46**	13.99	.92	14.64	1.93
Uncontrolled hostility	5.93	.20	6.73	.27*	5.80	.52	5.12	.34*
Anxiety/depression	8.93	.44	9.02	.44	9.48	.68	9.28	.69
	NAVIGATE							
	Non-Hispanic white (REF)		Non-Hispanic Black		Hispanic		Others	
	M	SD	M	SD	M	SD	M	SD
PANSS ^a								
Total score	63.61	1.45	68.20	2.05	66.34	2.26	73.13	4.49*
Negative symptoms	16.59	.56	17.53	.80	17.93	.95	19.01	1.31
Positive symptoms	19.60	.61	21.00	.78	19.39	.81	21.76	1.47
Disorganized thought	13.44	.38	14.23	.51	15.48	.54**	15.04	1.12
Uncontrolled hostility	6.02	.21	6.26	.35	6.19	.29	6.39	.57
Anxiety/depression	7.89	.31	9.17	.57*	7.40	.49	10.87	1.02**

^a Positive and Negative Syndrome Scale. Possible scores range from 30 to 120 with higher scores indicating greater psychosis symptom severity.

* $p < .05$, ** $p < .01$, *** $p < .001$;

Unadjusted percentage of participant engagement in key services for treatment conditions across the 24-month treatment period by race and ethnicity.



*** p<.001, ** p<.01, *p<.05