

Supplemental Table 1. Descriptive statistics by site

Variable	Site 1		Site 2		<i>t</i>	<i>df</i>	<i>p</i>	Number missing
	(N = 53)		(N = 68)					
Clinician variables	Mean	SD	Mean	SD				
Age	45.5	13.9	40.4	12.2	0.9	120	.377	0
	N	%	N	%	χ^2	<i>df</i>	<i>p</i>	
Sex, female	46	85	48	71	3.6	119	.056	0
Race, white	53	98	51	76	12.0	118	.001	1
Job role, case manager or therapist ^a	31	57	37	54	0.1	115	0.740	4
Clinician employed during the study (did not turnover)	38	72	52	77	0.6	119	.458	0
	(N = 116)		(N = 212)					
Client Variables	Mean	SD	Mean	SD	<i>t</i>	<i>df</i>	<i>p</i>	
Age	42.9	13.9	45.9	12.5	-2.0	337	.042	0
Familiarity ^b	16.1	27.9	29.2	48.4	-2.7	334	.007	3
Patient activation ^c	60.1	13.7	62.2	15.5	-1.2	336	.235	1
Physical health functioning ^d	41.2	7.9	42.7	8.2	-1.6	337	.107	0
Mental health functioning ^e	42.5	6.9	42.3	7.6	0.2	337	.806	0
Depression ^f	9.6	6.4	8.4	6.2	1.6	336	.113	1
Anxiety ^g	9.1	5.9	7.3	6.3	2.7	336	.008	1
	N	%	N	%	χ^2	<i>df</i>	<i>p</i>	
Sex, female	65	56	93	44	5.2	326	.022	0
Race, white	110	95	62	29	134.9	326	.000	0
Diagnosis ^h , SMI	48	43	166	79	48.0	326	.000	13

^a Job roles included ACT team member, medication provider, case manager, and therapist; note only 3 provided medication.

^b Client-Clinician Familiarity (months)

^c PAM-MH Scale: 0 – 100, with higher scores indicating greater patient activation

^d SF-12 Physical Health Scale: 0 – 100, with higher scores indicating better functioning

^e SF-12 Mental Health Scale: 0 – 100, with higher scores indicating better functioning

^f PHQ-9: Sum of 9 items, Scale: 0 (not at all) – 3 (nearly every day)

^g GAD-7: Sum of 7 items, Scale: 0 (not at all) – 3 (nearly every day)

^h SMI: bipolar disorder, schizophrenia spectrum disorders (schizophrenia was the most common diagnosis)

* $p < .05$; ** $p < .01$.