

Online supplement for DOI 10.1176/appi.ps.202100162

**Table 1A.** Participant Characteristics by Treatment Group

Characteristic & Variable	PHRP (N=32)	TAU (N=32)	$\chi^2$	<i>t</i>	<i>p</i>
Gender			7.23 <sup>a, c</sup>		.01
<i>Female</i> , freq (%)	27 (84.4)	17 (53.1)			
<i>Male</i> , freq (%)	5 (15.6)	15 (46.9)			
Ethnicity			1.31 <sup>b</sup>		.52
<i>Black or African-American</i> , freq (%)	3 (9.4)	4 (12.5)			
<i>White</i> , freq (%)	28 (87.5)	25 (78.1)			
<i>Hispanic or Latino</i> , freq (%)	1 (3.1)	3 (9.4)			
Marital status			1.36 <sup>b</sup>		.51
<i>Married</i> , freq (%)	3 (9.4)	2 (6.3)			
<i>Separated or Divorced</i> , freq (%)	3 (9.4)	1 (3.1)			
<i>Never Married</i> , freq (%)	26 (81.3)	29 (90.6)			
Employment			4.80 <sup>b</sup>		.09
<i>Unemployed</i> , freq (%)	8 (25.0)	16 (50.0)			
<i>Part-Time</i> , freq (%)	5 (15.6)	5 (15.6)			
<i>Full-Time</i> , freq (%)	19 (59.4)	11 (34.4)			
Inpatient Discharge Diagnosis					
<i>Trauma</i> , freq (%)	13 (40.6)	12 (37.5)	.07 <sup>a</sup>		.80
<i>Substance Use Disorder</i> , freq (%)	12 (37.5)	12 (37.5)	.00 <sup>a</sup>		1.00
<i>Depression</i> , freq (%)	24 (75)	26 (81.3)	.37 <sup>a</sup>		.55
<i>Anxiety</i> , freq (%)	13 (40.6)	14 (43.8)	.06 <sup>a</sup>		.80
<i>Bipolar</i> , freq (%)	4 (12.5)	1 (3.1)	1.95 <sup>a</sup>		.16
<i>Borderline</i> , freq (%)			7.05 <sup>b, c</sup>		.03
<i>Traits</i> , freq (%)	6 (18.8)	5 (15.6)			
<i>Diagnosis</i> , freq (%)	19 (59.3)	10 (31.3)			

Family History of Suicide Attempts			3.19 <sup>b</sup>	.20
<i>No Family History</i>	25 (78.1)	22 (68.8)		
<i>History in 1<sup>st</sup>/2<sup>nd</sup> Degree Relatives, freq (%)</i>	7 (21.9)	7 (21.9)		
<i>No Information in Record, freq (%)</i>	0 (0)	3 (9.4)		
Lethality of Prior Suicide Attempts			3.20 <sup>b</sup>	.20
<i>No Prior Suicide Attempts, freq (%)</i>	20 (62.5)	26 (81.3)		
<i>Low Lethality, freq (%)</i>	4 (12.5)	3 (9.4)		
<i>High Lethality, freq (%)</i>	8 (25.0)	3 (9.4)		
Suicide Attempt within 90 days of Admission	12 (37.5)	8 (25.0)	1.16 <sup>a</sup>	.28
Family Support			9.16 <sup>b, c</sup>	.01
<i>No Support, freq (%)</i>	2 (6.3)	12 (37.5)		
<i>Some Support, freq (%)</i>	22 (68.8)	15 (46.9)		
<i>Excellent Support, freq (%)</i>	8 (25.0)	5 (15.6)		
Age, mean ( <i>SD</i> )	25.4 (6.5)	28.1 (7.3)	-1.57 <sup>a</sup>	.08
Inpatient Length of Stay, mean ( <i>SD</i> )	6.97 (5.13)	6.72 (3.65)	.23 <sup>a</sup>	.82
Prior Suicide Attempts, mean ( <i>SD</i> )	1.78 (1.04)	1.22 (1.13)	2.07 <sup>a, c</sup>	.04
PHQ-9 Full Scale	20.81 (4.13)	15.00 (5.98)	4.52 <sup>a, c</sup>	<.001
Item 9: Suicidality	2.03 (.93)	1.28 (1.02)	3.06 <sup>a, c</sup>	.003

**Note.** PHQ-9 = Patient Health Questionnaire-9, Item 9: Thoughts that you would be better off dead, or of hurting yourself. <sup>a</sup> df=1. <sup>b</sup> df=2. <sup>c</sup> Included in multivariate analyses of predictors of rehospitalization.