

Table A. International Classification of Diseases (ICD) 10 Codes for Depression and Anxiety

ICD-10	Number of patients
Depression Diagnosis	
F32	17,838
F33	6,545
F34	988
F41	9,213
F53	265
Anxiety Diagnosis	
F06	18
F40	15
F41	43,604
F43	8,312

Figure A. Timeline of when each group of practices transitioned from pre-implementation to post-implementation status

Group	2016				2017				2018				2019				2020
	Qtr 1	Qtr 2	Qtr 3	Qtr 4	Qtr 1	Qtr 2	Qtr 3	Qtr 4	Qtr 1	Qtr 2	Qtr 3	Qtr 4	Qtr 1	Qtr 2	Qtr 3	Qtr 4	Qtr 1
1	Light	Dark															
2	Light	Dark															
3	Light	Dark															
4	Light	Dark															
5	Light	Dark															
6	Light	Dark	Dark	Dark	Dark	Dark	Dark										
7	Light	Dark	Dark	Dark	Dark	Dark											

Qtr = quarter

Light grey = pre-implementation period

Dark grey = post-implementation period

Table B. The Adjusted Odds of Having a Behavioral Health Visit

Adjusted odds of having a behavioral visit									
	Depression Only			Anxiety Only			Depression and Anxiety		
	AOR	95%CI		AOR	95%CI		AOR	95%CI	
Post period (versus pre-period)	4.35	3.50	5.41	4.27	3.57	5.11	3.26	2.77	3.84
Increase in 1 year of age	0.97	0.97	0.98	0.98	0.98	0.98	0.98	0.98	0.99
Female (versus male)	1.21	1.06	1.39	1.14	1.04	1.25	1.31	1.18	1.45
Black race (versus White race)	1.52	1.27	1.82	1.48	1.30	1.70	1.89	1.64	2.18
Other race (versus White race)*	1.60	1.27	2.02	1.47	1.27	1.71	1.66	1.42	1.94
Medicare (versus private)	1.06	0.85	1.32	0.95	0.79	1.13	0.97	0.81	1.17
Medicaid (versus private)	0.84	0.70	1.02	1.04	0.90	1.19	0.94	0.82	1.08
Other (versus private)	1.05	0.69	1.61	0.90	0.66	1.22	1.16	0.84	1.60
Married or partnered (versus not)	0.71	0.61	0.82	0.89	0.80	0.98	0.89	0.80	0.99

The other race category was composed of 55% of individuals who did not identify their race, 30% who identified as Multiracial/Multicultural, 15% who identified as Asian, and 2% who identified as American Indian/Alaskan Native.