

On-Line Supplemental Table: Change in APS Participants' Reported Confidence Before and After Completing Training

	Pre n	%	Post n	%	P
Identify suicide warning signs in a patient. (n=873)					<0.01
Not confident/ not very confident	158	18	3	0	
Neutral	183	21	25	3	
Extremely confident/ somewhat confident	532	61	845	97	
Know what to do if a PHQ-9 score is high. (n=872)					<0.01
Not confident/ not very confident	310	36	9	1	
Neutral	103	12	36	4	
Extremely confident/ somewhat confident	459	53	827	95	
Write a safety plan with a patient. (n=872)					<0.01
Not confident/ not very confident	390	45	37	4	
Neutral	166	19	105	12	
Extremely confident/ somewhat confident	316	36	730	83	
Listen without judgment to a patient. (n=868)					<0.01
Not confident/ not very confident	22	3	0	0	
Neutral	52	6	12	1	
Extremely confident/ somewhat confident	794	91	856	99	
Ask a patient directly about suicide. (n=868)					<0.01
Not confident/ not very confident	65	8	4	0	
Neutral	73	8	16	2	
Extremely confident/ somewhat confident	730	84	848	98	
Connect a patient to additional resources. (n=864)					<0.01
Not confident/ not very confident	179	21	4	0	
Neutral	131	15	38	5	
Extremely confident/ somewhat confident	554	64	822	95	
Deliver a message to all patients about med safety. (n=871)					<0.01
Not confident/ not very confident	125	14	8	1	
Neutral	177	20	42	5	
Extremely confident/ somewhat confident	569	66	821	94	
Deliver a message to all patients about firearm safety. (n=865)					<0.01
Not confident/ not very confident	140	16	10	1	
Neutral	171	20	49	6	
Extremely confident/ somewhat confident	554	64	806	93	