

Table 1. Characteristics of respondents in the Collaborative Psychiatric Epidemiology Survey, by percentage reporting psychotic symptoms

Characteristic	12-month prevalence of any psychotic symptoms		
	%	95% CI	p
Race			<.001
Asian	1.8	1.2-2.8	
Latino	3.4	2.6-4.5	
Black	4.5	3.7-5.4	
White	2.2	1.6-3.0	
Other	0.3	.1-2.0	
Sex			0.56
Male	2.7	2.0-3.5	
Female	3.0	2.4-3.6	
Age			0.03
18-25	3.6	2.5-5.1	
26-45	3.4	2.7-4.4	
46-65	2.2	1.5-3.2	
>65	1.7	.9-3.0	
Marital Status			<.001
Married	1.7	1.3-2.3	
Not Married	4.4	3.5-5.4	
Education (years)			0.40
0-11	3.5	2.5-4.8	
12	2.5	1.8-3.3	
13-15	2.7	2.1-3.6	
>=16	2.8	1.9-4.0	
Income			<.001
<\$22,500	4.2	3.3-5.3	
>=\$22,500	2.3	1.9-2.9	
Region			0.05
Northeast	2.1	1.4-3.2	
Midwest	3.6	2.5-5.1	
South	2.3	1.9-2.9	
West	3.4	2.6-4.6	

N=11,615. Data were adjusted for weighting and design effects. All data were not available for all subjects.

Table 2. Psychological factors of respondents in the Collaborative Psychiatric Epidemiology Surveys, by percentage reporting psychotic symptoms

**12-month prevalence of any
psychotic symptoms**

Psychological factors	%	95% CI	p
Higher distress			<.001
No	1.2	.8-1.8	
Yes	4.3	3.7-5.0	
Dysfunction days present in past 30 days			0.03
0-2 days	2.3	1.8-2.9	
3-30 days	3.8	2.5-5.8	
Lifetime substance use disorder			<.001
No	2.5	2.1-3.0	
Yes	5.2	3.7-7.2	
Lifetime posttraumatic stress disorder			<.001
No	2.5	2.1-3.0	
Yes	7.6	5.6-10.2	

N=11,615. Data were adjusted for weighting and design effects. All data were not available for all subjects.

Table 3. Logistic regression analysis of characteristics associated with 12-month prevalence of psychotic symptoms among respondents in the Collaborative Psychiatric Epidemiology Surveys

Characteristic	12-month prevalence	
	OR	95% CI
Race (reference: white)		
Asian	0.9	.5-1.7
Latino	1.7	1.004-2.8
Black	3.5	1.6-7.4
Other	0.1	0-0.5
Female (reference: male)	0.8	.5-1.2
Age (reference: >65)		
18-25	1.4	.6-3.5
26-45	1.9	.7-5.3
46-65	1.2	.4-3.3
Not Married (reference: married)	2.5	1.5-4.1
Education (reference: ≥ 16)		
0-11 years	0.8	.3-2.0
12 years	0.8	.4-1.4
13-15 years	0.8	.4-1.5
Income Level <\$22,500 (reference: ≥ \$22,500)	1.2	.7-2.1
Region (reference: West)		
Northeast	0.5	.3-1.0
Midwest	0.9	.5-1.7
South	0.6	.4-0.9
Lifetimes substance use disorder (reference: no)	1.3	.8-2.0
Lifetime posttraumatic stress disorder (reference: no)	1.8	1.1-3.2
Higher Distress (reference: no)	3.0	1.7-5.2
Dysfunction on 3-30 days in past 30 days (reference: 0-2 days)	1.2	.8-2.0

N=11,615. Data were adjusted for weighting and design effects. All data were not available for all subjects.