

Supplemental Table. Unadjusted Associations among WLQ Subscale Scores, Work Productivity Loss, and Work Productivity Cost and Sociodemographic/Background Characteristics and Mental Health/Substance Use Assessment Outcomes (n=473).

Sociodemographic/ Background Characteristics	Work Limitations Questionnaire Subscale					Productivity Loss
	Time Management	Physical Demands	Mental/ Interpersonal Demands	Output Demands		
Employment Type						
Full time	32.7 \pm 32.1	34.0 \pm 33.4	27.7 \pm 27.4	17.9 \pm 27.81	6.9 \pm 6.1	
Part time	29.4 \pm 33.3	33.9 \pm 35.2	26.6 \pm 29.8	20.2 \pm 28.52	6.8 \pm 6.4	
	$F_{(1,471)}$, <i>p</i> -value	.75, .39	.001, .98	.11, .75	.48, .49	.003, .96
Age Group						
18-29 years	30.1 \pm 31.4	31.2 \pm 32.4	25.8 \pm 27.2	17.0 \pm 27.37	6.4 \pm 5.9	
30-39 years	35.6 \pm 32.7	38.2 \pm 36.1	29.8 \pm 27.2	17.7 \pm 28.13	7.3 \pm 6.3	
40 years old and over	32.0 \pm 33.3	34.4 \pm 33.3	28.0 \pm 29.3	20.9 \pm 28.61	7.2 \pm 6.4	
	$F_{(2,470)}$, <i>p</i> -value	1.12, .33	1.72, .18	.81, .45	.84, .43	1.00, .37
Gender						
Female	39.5 \pm 31.0	36.2 \pm 34.9	31.8 \pm 28.4	27.2 \pm 29.9	8.5 \pm 6.4	
Male	31.1 \pm 32.4	33.7 \pm 33.6	26.9 \pm 27.7	17.1 \pm 27.5	6.6 \pm 6.1	
	$F_{(1,471)}$, <i>p</i> -value	3.41, .07	.28, .60	1.56, .21	6.63, .01	4.63, .03
Relationship Status						
Married/Partnered	34.8 \pm 33.5	37.9 \pm 35.33	28.3 \pm 28.89	18.6 \pm 28.0	7.2 \pm 6.3	
Single	29.9 \pm 31.3	30.9 \pm 32.1	26.9 \pm 27.0	18.1 \pm 27.9	6.6 \pm 6.0	
	$F_{(1,471)}$, <i>p</i> -value	2.72, .10	5.09, .02	.30, .58	.03, .86	1.13, .29
Ethnicity						
Other	34.2 \pm 32.5	38.3 \pm 34.1	30.0 \pm 27.8	21.3 \pm 29.1	7.6 \pm 6.3	
Non-Hispanic White	30.9 \pm 32.2	31.4 \pm 33.3	26.1 \pm 27.2	16.6 \pm 27.1	6.4 \pm 6.0	
	$F_{(1,471)}$, <i>p</i> -value	1.18, .28	4.58, .03	2.18, .14	3.24, .07	4.04, .05
Financial Situation						
Have at least enough to get by	31.2 \pm 32.0	32.9 \pm 33.7	26.1 \pm 26.8	17.4 \pm 27.3	6.6 \pm 6.0	
Can't make ends meet	39.1 \pm 35.6	42.1 \pm 31.7	37.7 \pm 33.2	25.0 \pm 31.5	9.0 \pm 6.8	
	$F_{(1,471)}$, <i>p</i> -value	2.94, .09	3.60, .06	8.58, .004	3.59, .06	7.28, .01
Significant Head Injury						
No	28.2 \pm 31.3	30.3 \pm 33.2	24.4 \pm 26.2	16.8 \pm 26.5	6.2 \pm 5.9	
Yes	37.5 \pm 33.0	39.5 \pm 33.7	31.5 \pm 29.5	19.9 \pm 29.4	7.8 \pm 6.4	
	$F_{(1,466)}$, <i>p</i> -value	9.45, .002	8.50, .004	7.55, .01	1.41, .24	8.02, .01
Taken Medication for Depression, Anxiety, or Nerves in Prior Month						
No	30.2 \pm 32.5	32.3 \pm 33.7	26.5 \pm 28.2	17.0 \pm 27.8	6.5 \pm 6.1	
Yes	37.0 \pm 31.4	38.3 \pm 33.5	30.2 \pm 26.7	21.8 \pm 28.2	7.8 \pm 6.1	
	$F_{(1,471)}$, <i>p</i> -value	4.27, .04	2.97, .09	1.67, .20	2.76, .10	4.31, .04
SF-12 Mental Component Score	<i>r</i> , <i>p</i> -value	-.45, <.001	-.23, <.001	-.51, <.001	-.37, <.001	-.51, <.001
SF-12 Physical Component Score	<i>r</i> , <i>p</i> -value	-.22, <.001	-.16, <.001	-.11, .02	-.21, <.001	-.21, <.001
Mental Health/Substance						

Use Assessment Outcomes	Work Limitations Questionnaire Subscale					
	Time Management	Physical Demands	Interpersonal Demands	Output Demands	Mental/	Productivity Loss
					Productivity Loss	
Depressive Symptom Severity ± PHQ-9	<i>r, p-value</i>	.44, <.001	.24, <.001	.50, <.001	.37, <.001	.51, <.001
Major Depression						
No		25.3 ± 29.6	29.9 ± 33.7	19.9 ± 23.5	13.2 ± 24.2	5.3 ± 5.3
Yes		47.4 ± 33.0	43.1 ± 31.9	44.6 ± 29.7	29.8 ± 32.2	10.4 ± 6.4
	$F_{(1,471)}, p-value$	52.18, <.001	15.87, <.001	94.51, <.001	38.36, <.001	82.15, <.001
Suicide Ideation Severity						
Non-High Risk		30.0 ± 32.0	33.2 ± 33.9	25.7 ± 26.9	16.2 ± 26.4	6.4 ± 5.9
High Risk		43.2 ± 31.7	37.8 ± 32.8	37.0 ± 30.9	29.7 ± 32.9	9.4 ± 7.0
	$F_{(1,471)}, p-value$	10.68, .001	1.16, .28	10.46, .001	15.10, <.001	14.96, <.001
PTSD Symptom Severity ± PCL	<i>r, p-value</i>	.40, <.001	.22, <.001	.41, <.001	.35, <.001	.45, <.001
PTSD						
No		23.0 ± 29.1	29.0 ± 33.6	19.4 ± 24.6	11.8 ± 24.3	5.0 ± 5.3
Yes		43.2 ± 32.6	40.1 ± 33.0	37.5 ± 28.4	26.4 ± 30.0	9.2 ± 6.3
	$F_{(1,471)}, p-value$	50.42, <.001	13.05, <.001	55.57, <.001	34.10, <.001	63.45, <.001
Generalized Anxiety Disorder &/or Panic						
No		21.3 ± 26.9	28.8 ± 33.8	18.6 ± 24.3	11.2 ± 23.30	4.8 ± 5.1
Yes		46.0 ± 33.5	40.6 ± 32.6	38.9 ± 28.0	27.5 ± 30.63	9.6 ± 6.3
	$F_{(1,471)}, p-value$	79.05, <.001	14.88, <.001	71.06, <.001	43.16, <.001	84.41, <.001
Binges in Prior 3 Months	<i>r, p-value</i>	.11, .01	.10, .03	.14, .002	.14, .002	.16, <.001
Average Number of Alcoholic Drinks in Past Week	<i>r, p-value</i>	.07, .16	.08, .08	.09, .5	.07, .11	.10, .04
Alcohol Dependent &/or Illicit Drug Use						
No		30.3 ± 31.8	31.7 ± 33.6	26.0 ± 27.4	16.3 ± 26.2	6.4 ± 6.0
Yes		39.9 ± 33.6	43.8 ± 32.8	34.1 ± 28.9	27.0 ± 33.4	8.9 ± 6.5
	$F_{(1,471)}, p-value$	6.43, .01	9.28, .002	6.17, .01	10.65, .001	12.60, <.001

Note. Cell values represent estimates (means and standard deviations (M (SD)) or, where denoted, correlation coefficients (*r*) from unadjusted, bivariate analyses of the association among sociodemographic, clinical, and psychiatric factors and WLQ subscales. Tests of significance included one-way ANOVA and Pearson correlations for categorical and continuous variables, respectively. Analyses were two-tailed. Continuous measures included the following: PHQ-9= Patient Health Questionnaire (range= 0-27, higher scores indicate greater symptomatology); PCL= Post-Traumatic Stress Disorder (PTSD) Patient Checklist (range=17-85, higher scores indicate greater symptomatology); SF-12 = Short Form Health Survey of the Medical Outcomes Study (range=0-100, higher scores indicate greater mental/physical functioning); Binges in prior 3 months (range=0-150); Average number of alcoholic drinks in past week (range=0-29). WLQ Subscale range (0-100) with the exception of Productivity Loss Subscale (range=0-25) and Productivity Cost (range=0-8116).