













































































































































































































































457. Raskino C, Pearson DA, Baker CJ, Lifschitz MH, O'Donnell K, Mintz M, Nozyce M, Bruwors P, McKinney RE, Jimenez E, England JA (Pediatric AIDS Clinical Trials Group 152 Study Team): Neurologic, neurocognitive, and brain growth outcomes in human immunodeficiency virus-infected children receiving different nucleoside antiretroviral regimens. *Pediatrics* 1999; 104(3):e32 [A]
458. Tepper VJ, Farley JJ, Rothman MI, Houck DL, Davis KF, Collins-Jones TL, Wachtel RC: Neurodevelopmental/neuroradiologic recovery of a child infected with HIV after treatment with combination antiretroviral therapy using the HIV-specific protease inhibitor ritonavir. *Pediatrics* 1998; 101:e7 [G]
459. Cesena M, Lee D, Cebollero A, Steingard R: Case study: behavioral symptoms of pediatric HIV-1 encephalopathy successfully treated with clonidine. *J Am Acad Child Adolesc Psychiatry* 1995; 34:302–306 [G]
460. Anand A, Carmosino L, Glatt A: Management of recalcitrant pain in a pediatric acquired immunodeficiency syndrome patient. *Pediatr Infect Dis J* 1993; 12:159–160 [G]
461. Hirschfeld S, Moss H, Dragisic K, Smith W, Pizzo PA: Pain in pediatric human immunodeficiency virus infection: incidence and characteristics in a single-institution pilot study. *Pediatrics* 1996; 98:449–452 [G]
462. Grinspoon S, Corcoran C, Askari H, Schoenfeld D, Wolf L, Burrows B, Walsh M, Hayden D, Parlman K, Anderson E, Basgoz N, Klibanski A: Effects of androgen administration in men with the AIDS wasting syndrome: a randomized, double-blind, placebo-controlled trial. *Ann Intern Med* 1998; 129:18–26 [A]
463. Rabkin JG, Rabkin R, Wagner GJ: Testosterone treatment of clinical hypogonadism in patients with HIV/AIDS. *Int J STD AIDS* 1997; 8:537–545 [F]
464. Hengge UR, Baumann M, Maleba R, Brockmeyer NH, Goos M: Oxymetholone promotes weight gain in patients with advanced human immunodeficiency virus (HIV-1) infection. *Br J Nutr* 1996; 75:129–138 [A]
465. Miller K, Corcoran C, Armstrong C, Caramelli K, Anderson E, Cotton D, Basgoz N, Hirschhorn L, Tuomala R, Schoenfeld D, Daugherty C, Mazer N, Grinspoon S: Transdermal testosterone administration in women with acquired immunodeficiency syndrome wasting: a pilot study. *J Clin Endocrinol Metab* 1998; 83:2717–2725 [B]
466. Wagner G, Rabkin J, Rabkin R: Exercise as a mediator of psychological and nutritional effects of testosterone therapy in HIV+ men. *Med Sci Sports Exerc* 1998; 30:811–817 [B]
467. Strawford A, Barbieri T, Van Loan M, Parks E, Catlin D, Barton N, Neese R, Christiansen M, King J, Hellerstein MK: Resistance exercise and supraphysiologic androgen therapy in eugonadal men with HIV-related weight loss: a randomized controlled trial. *JAMA* 1999; 281:1282–1290 [A]
468. Breitbart WS, Rosenfeld B, Kaim M, Funesti-Esch J: Psychostimulants for fatigue in the HIV+ patient: preliminary findings of a placebo-controlled trial of methylphenidate vs pemoline. *Psychosomatics* 1999; 40:160–161 [A]
469. Rabkin JG, Ferrando SJ, Wagner GJ, Rabkin R: DHEA treatment for HIV+ patients: effects on mood, androgenic and anabolic parameters. *Psychoneuroendocrinology* 2000; 25:53–68 [B]
470. Holzemer WL, Henry SB, Reilly CA: Assessing and managing pain in AIDS care: the patient perspective. *J Assoc Nurses AIDS Care* 1998; 9:22–30 [G]
471. Kemper CA, Kent G, Burton S, Deresinski SC: Mexiletine for HIV-infected patients with painful peripheral neuropathy: a double-blind, placebo-controlled, crossover treatment trial. *J Acquir Immune Defic Syndr Hum Retrovirol* 1998; 19:367–372 [A]
472. Shlay JC, Chaloner K, Max MB, Flaws B, Reichelderfer P, Wentworth D, Hillman S, Brizz B, Cohn DL: Acupuncture and amitriptyline for pain due to HIV-related peripheral neuropathy: a randomized controlled trial: Terry Beinr Community Programs for Clinical Research on AIDS. *JAMA* 1998; 280:1590–1595 [A]
473. Hall MC, Ahmad S: Interaction between sildenafil and HIV-1 combination therapy (letter). *Lancet* 1999; 353:2071–2072 [G]